

abinvasives.ca info@abinvasives.ca



Last Updated January 2014



Hybrid Knapweed

Centaurea x psammogena



www.easterncoloradowildflowers.com

ww.easterncoloradowildflowers.com

Overview:

Hybrid knapweed is a cross of diffuse (*C.diffusa*) and spotted (*C. stoebe*) knapweed. However, among the literature there are various accounts about which subspecies of spotted knapweed was involved, and the taxonomy of spotted knapweed varies around the world as well. There is also disagreement regarding whether the hybridization occurred before import to North America and whether it still occurs among alien, naturalized populations. *Centaurea x psammogena* was first described in 1909 by Gyula Gáyer.

Like the many other knapweeds, it is a short-lived perennial forb of the Aster family. Its origin is southwestern Europe and method of introduction is speculative. The morphology of hybrid knapweed is extremely variable; it may be intermediate, or may closely resemble one or the other of the parent species.²

Habitat:

Hybrid knapweed grows in sandy, coarse or loamy, well-drained soils, moist but not wet,

and generally in full sun.

Identification:

Stems: Are single to several, erect, branched in the upper stem, and growing 80-150 cm tall. Stems may be finely hairy.

Leaves: Basal and lower stems leaves are 10-20 cm long and have short petioles. The leaf margins may be deeply lobed or bi-pinnately divided. Leaves decrease in size on upper stem, are sessile (no petioles), and may be simple or lobed. All leaves may be minutely hairy.

Flowers: Are borne singly at the ends of stems and composed of 20-40 florets which may be purple colored (rarely white) or have white central florets with a purple outer ring of florets.¹ Flower bases are covered with bracts which are brown to black (rarely straw-coloured), and spines may be absent or short.² Seeds always bear a pappus and radiant heads.²

Prevention:

Hybrid knapweed may be available through

the horticultural trade. Other methods of spread would be through seed contaminated soil, or plants being caught in vehicles.

Control:

Grazing: Unknown. Invasive plants should never be considered as forage.

Mechanical: Small infestations may be hand-pulled or dug before seed set, taking to care to remove the entire root.

Chemical: Currently no selective herbicides are registered for use on hybrid knapweed. Always check product labels to ensure the herbicide is registered for use on the target plant in Canada by the Pest Management Regulatory Agency. Always read and follow label directions. Consult your local Agricultural Fieldman or Certified Pesticide Dispenser for more information.

Biological: None researched to date.

REFERENCES

- 1 MacBryde, Bruce. A Baker's Dozen Worst Weeds Invading the Estes Valley. European Hybrid Knapweed. Preliminary Edition 2011.
- 2 Centaurea diffusa in Flora of North America. www.efloras.org

